

Children's dosing for pain or fever



Kids deserve the best.












Liquid medicine for children

Acetaminophen

liquid medicine 160mg/5mL
(Tylenol) give by mouth

Ibuprofen

liquid medicine 100mg/5mL
(Motrin or Advil) give by mouth

Directions	Give every 4-6 hours as needed for fever or pain. Do not give more than 5 times in a 24 hours period.	Visual shows use of 10 mL syringe. Be sure to check your syringe size.	Give every 6 hours as needed for fever or pain.
4-8 pounds	1mL		Do not give to infants less than 6 months unless told by child's doctor.
8-10 pounds	2 mL		2 mL
11-16 pounds	3 mL		3 mL
15-20 pounds	4 mL		4 mL
21-25 pounds	5 mL		5 mL
26-29 pounds	6 mL		6 mL
30-39 pounds	8 mL		8 mL
40-49 pounds	10 mL		10 mL
50-59 pounds	12 mL		12 mL
60-79 pounds	15 mL		15 mL
80 pounds+	20 mL		20 mL

Children's dosing for pain or fever



Kids deserve the best.

Tablets for children

	Acetaminophen (Tylenol)				Ibuprofen (Motrin or Advil)		
Directions	Give every 4-6 hours as needed for fever or pain. Do not give more than 5 times in a 24 hours period.				Give every 6 hours as needed for fever or pain.		
Milligrams (mg)	Chewable 80 mg	Chewable 160 mg	Adult tabs 325 mg	Adult tabs 500 mg	Chewable 50 mg	Chewable 100 mg	Adult tabs 200 mg
12-17 pounds	1 tab				1 tab		
18-23 pounds	1½ tabs				1 tab		
24-35 pounds	2 tabs	1 tab			2 tabs	1 tab	
36-47 pounds	3 tabs	1½ tabs			3 tabs	1½ tabs	
48-59 pounds	4 tabs	2 tabs	1 tab		4 tabs	2 tabs	1 tab
60-71 pounds	5 tabs	2½ tabs	1 tab		5 tabs	2½ tabs	1 tab
72-95 pounds	6 tabs	3 tabs	1½ tabs	1 tab	6 tabs	3 tabs	1½ tabs
96+ pounds	8 tabs	4 tabs	2 tabs	1 tab	7 tabs	4 tabs	2 tabs